

* $p < 0.05$; ** $p < 0.01$.




Figure 2 Correlation between 24 hour urinary excretion of creatinine and uric acid in (A) patients with primary gout; (B) normal male subjects.

Discussion

In this study significant positive correlation between 24 hour urinary creatinine and uric acid excretion was shown. In the steady state urinary excretion of uric acid over 24 hours is thought to represent the urate production over the same period.¹⁷ Although 24 hour urinary excretion of creatinine may be affected by the dietary creatine content, it seems that there is a close association between creatinine and uric acid synthesis.

One possible mechanism for this association is the increased supply of creatine from the diet. This exogenous creatine is subsequently metabolised to creatine phosphate, then creatinine. The phosphate donor is ATP. Increased degradation of ATP has been shown to cause accelerated urate synthesis.⁷⁻¹⁴

Another possible mechanism for the association between creatinine and uric acid synthesis is endogenous creatine synthesis, by methylation of guanidinoacetic acid. In this reaction S-

adenosylmethionine is converted to S-adenosylhomocysteine, which is then catalysed to adenosine. Accelerated creatine synthesis may cause increased synthesis of adenosine, which is degraded subsequently to inosine, hypoxanthine, xanthine, and uric acid. Fasting subjects excreted 400 to 900 mg of creatinine in 24 hours, indicating endogenous creatine synthesis (data not shown). It seems, therefore, that large amounts of adenosine are produced by methylation.

In addition, patients with gout had increased 24 hour urinary creatinine and uric acid excretion. Although this study did not measure either endogenous creatinine or uric acid synthesis directly, it is possible that the hyperuricaemia seen in some patients with primary gout is due to accelerated endogenous creatinine synthesis.

We are carrying out further studies of the relation between creatinine and uric acid metabolism in animals.

- 1 Sperling O, Eilam G, Persky-Brosh S, de Vries A. Accelerated erythrocyte 5-phosphoribosyl-1-pyrophosphate synthesis. A familial abnormality associated with excessive uric acid production and gout. *Biochem Med* 1972; 6: 310-6.
- 2 Becker M A, Meyer L J, Seegmiller J E. Gout with purine overproduction due to increased phosphoribosylpyrophosphate synthetase activity. *Am J Med* 1973; 55: 232-42.
- 3 Lesch M, Nyhan W L. A familial disorder of uric acid metabolism and central nervous system function. *Am J Med* 1964; 36: 561-70.
- 4 Seegmiller J E, Rosenbloom F M, Kelley W N. Enzyme defect associated with a sex-linked human neurological disorder and excessive purine synthesis. *Science* 1967; 155: 1682-4.
- 5 Kelley W N, Rosenbloom F M, Henderson J F, Seegmiller J E. A specific enzyme defect in gout associated with overproduction of uric acid. *Proc Natl Acad Sci USA* 1967; 57: 1735-9.
- 6 Morton N E, Lalouel J M. Genetic epidemiology of Lesch-Nyhan disease. *Am J Hum Genet* 1977; 29: 304-7.
- 7 Fox I H. Adenosine triphosphate degradation in specific disease. *J Lab Clin Med* 1985; 106: 101-10.
- 8 Fox I H. Metabolic basis for disorders of purine nucleotide degradation. *Metabolism* 1981; 30: 616-34.
- 9 Fox I H, Kelly W N. Studies on the mechanism of fructose induced hyperuricemia in man. *Metabolism* 1972; 21: 713-21.
- 10 Faller J, Fox I H. Ethanol-induced hyperuricemia. Evidence for increased urate production by activation of adenine nucleotide turnover. *N Engl J Med* 1982; 307: 1598-602.
- 11 Woolliscroft J O, Colfer H, Fox I H. Hyperuricemia in acute illness: a poor prognostic sign. *Am J Med* 1982; 72: 58-62.
- 12 Fox I H, Palella T D, Kelley W N. Hyperuricemia: a marker for cell energy crisis. *N Engl J Med* 1987; 317: 111-2.
- 13 Sutton J R, Toews C J, Ward G R, Fox I H. Purine metabolism during strenuous muscular exercise in man. *Metabolism* 1980; 29: 254-60.
- 14 Green H J, Fraser I G. Differential effects of exercise intensity on serum uric acid concentration. *Med Sci Sports Exerc* 1988; 20: 55-9.
- 15 Liddle L, Seegmiller J E, Laster L. The enzymatic spectrophotometric method for determination of uric acid. *J Lab Clin Med* 1959; 54: 903-13.
- 16 Bonsnes R W, Taussky H H. On the colorimetric determination of creatinine by the Jaffe reaction. *J Biol Chem* 1945; 158: 581-91.
- 17 Seegmiller J E, Grayzel A I, Laster L, et al. Studies of uric acid pool size and turnover rate. *Ann Rheum Dis* 1969; 28: 366-73.